

INSTRUCTIONS FOR USE
OTOVEL (OH-toe-vel)
(ciprofloxacin and fluocinolone acetonide)
otic solution


Read this Instructions for Use that comes with OTOVEL before you start using it and each time you get a refill. There may be new information. This information does not take the place of talking with your healthcare provider about your medical condition or treatment.

Important information about OTOVEL:


- **OTOVEL is for use in the ear only (otic use).** Do not inject OTOVEL or use OTOVEL in the eye.
- Use OTOVEL exactly as your healthcare provider tells you to use it.

How should I use OTOVEL?


Step 1. You or your caregiver should wash their hands with soap and water.




Step 2. Gently clean any fluid (discharge) from the outer ear using a clean cloth or tissue. **Do not** put a cotton swab or any other object in the ear canal.




Step 3. Remove OTOVEL from the protective foil pouch. Pull apart 1 single-dose vial of OTOVEL as shown, by tearing along the dotted lines (perforations) until it is fully separated.




Step 4. Warm the dose of OTOVEL by holding the vial in your hand for **1 to 2 minutes**.




Step 5. Twist off the vial cap in the direction of the arrow as shown.




Step 6. The person receiving OTOVEL should be on his/her side with the infected ear up as shown.



Step 7. Hold the vial of OTOVEL in your hand and place the vial close to the ear. Let the entire dose of OTOVEL fall into the affected ear.



Step 8. Gently press the part of the ear known as the **tragus 4 times** using a pumping motion as shown. This will allow the drops of OTOVEL to enter the middle ear.



Step 9. Remain on your side with the affected ear facing upward for **1 minute**.



Step 10. If your healthcare provider has told you to use OTOVEL in both ears, repeat Steps 2-9 for the other ear.

Step 11. Safely throw away OTOVEL vials after use.

This Instructions for Use has been approved by the Food and Drug Administration.